

The placenta and neurodevelopment: sex differences in prenatal vulnerability

Tracy L. Bale

To cite this article: Tracy L. Bale (2016) The placenta and neurodevelopment: sex differences in prenatal vulnerability, Dialogues in Clinical Neuroscience, 18:4, 459-464, DOI: [10.31887/DCNS.2016.18.4/tbale](https://doi.org/10.31887/DCNS.2016.18.4/tbale)

To link to this article: <https://doi.org/10.31887/DCNS.2016.18.4/tbale>



Copyright: © 2016 Institut la Conference Hippocrate - Servier Research Group



Published online: 01 Apr 2022.



Submit your article to this journal [↗](#)



Article views: 371



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 48 View citing articles [↗](#)

Brief report

The placenta and neurodevelopment: sex differences in prenatal vulnerability

Tracy L. Bale, PhD



Prenatal insults, such as maternal stress, are associated with an increased neurodevelopmental disease risk and impact males significantly more than females, including increased rates of autism, mental retardation, stuttering, dyslexia, and attention deficit/hyperactivity disorder (ADHD). Sex differences in the placenta, which begin with sex chromosomes, are likely to produce sex-specific transplacental signals to the developing brain. Our studies and others have identified X-linked genes that are expressed at higher levels in the female placenta. Through a genome-wide screen after maternal stress in mice, we identified the X-linked gene O-linked N-acetylglucosamine transferase (OGT) and demonstrated its causality in neurodevelopmental programming producing a male-specific stress phenotype. Elucidating the sex-specific molecular mechanisms involved in transplacental signals that impact brain development is key to understanding the sex bias in neurodevelopmental disorders and is expected to yield novel insight into disease risk and resilience.

© 2016, AICH – Servier Research Group

Dialogues Clin Neurosci. 2016;17:459–464.

Keywords: *early prenatal stress; O-linked N-acetylglucosamine transferase; maternal stress; neurodevelopmental disorder; transplacental signal*

Introduction

Males are four to eight times likelier than females to be affected by a neurodevelopmental disorder; this complex statistic is dependent on a number of variables, including comorbidities and genetic variants.^{1–10} It is clear that prenatal insults, such as maternal stress, are associated with increased neurodevelopmental disease risk and impact males significantly more than females, including increased rates of autism, mental retardation, stuttering, dyslexia, and attention deficit/hyperactivity disorder (ADHD).^{11–15} Suggested explanations as to why males present with higher rates of these disorders range from the male brain developing more slowly than the female—placing the male brain at increased vulnerability over a greater period of time—to the perception that males are less able to compensate when unilateral brain damage occurs developmentally.^{16–19} A more recent and exciting hypothesis focuses on the critical involvement of the placenta in relaying sex-specific transplacental signals to the developing brain regarding changes or perturbations in the maternal milieu.^{20–26}

Furthermore, such maternal insults during pregnancy

Author affiliations: Department of Biomedical Sciences, School of Veterinary Medicine and Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA

Address for correspondence: Tracy L. Bale, PhD, Professor of Neuroscience, 410F Hill Pavilion, University of Pennsylvania, 380 South University Ave, Philadelphia, Pennsylvania 19104 USA (email: tbale@vet.upenn.edu)

Brief report

produce robust sex differences in transcriptional responses in the placenta.^{22-24,27} This Brief Report focuses on the mechanistic evidence that the known sex bias in neurodevelopmental disorder risk begins with placental sex differences.

Surprisingly, placental sex (XX vs XY) is a major determinant in the magnitude and functional responses of the placenta to maternal perturbations during pregnancy, where the female placenta appears to be protective.^{7,20,25,28-43} Placental cellular mechanisms give rise to sex-specific neurodevelopmental changes and are expected to provide novel insight into disease risk and resilience. The placenta is a highly dynamic endocrine tissue, producing nutrients and growth factors and maintaining a protective barrier, all critical for appropriate fetal development.^{22,27,44-47} Sex differences in the placenta begin as early as the trophoctoderm stage and remain robust throughout trophoblast cell differentiation, all of which occurs in the first third of gestation in humans and rodents. Sex-chromosome genes are differentially expressed early on, suggesting a mechanism whereby sex differences could be driven by the ability of the placenta to respond to changes in the maternal milieu and ultimately by the transplacental signals received by the developing fetus. We and others have reported significant differences in male and female placental gene expression patterns across gestation in rodent models; these differences were similarly found in human placenta.^{11,20,23,24,39,48} Despite differences in placental structure between rodents and humans, such studies suggest that key functional similarities exist and that mechanisms uncovered in animal models are translatable.²⁴

Placental mechanisms

Why male, but not female, placentas are consistently so responsive to changes in the maternal environment (eg, in response to maternal stress, infection, and diet) remains a key question in defining potential sex-specific disease mechanisms.^{26,49-55} Addressing the possible mechanisms underlying these sex differences, animal models have demonstrated (through functional and causal studies) the importance of specific sex-chromosome genes. As an example, in our established mouse model of early prenatal stress (EPS), male, but not female, offspring present with increased stress sensitivity as adults, including increased hypothalamic-pituitary-

adrenal (HPA) stress axis activity, reduced growth after weaning, and hypothalamic mitochondrial dysfunction.^{34,35,39,56} In a genome-wide screen for sex-specific changes in the placenta after EPS, we identified the X-linked gene *O*-linked N-acetylglucosamine transferase (*OGT*) as a top candidate for having an important role in the placenta in guiding broad chromatin changes that are probably important in placental function because of its sex specificity in expression and response to maternal stress. In the placenta, *OGT* escapes X inactivation; consequently, levels are twice as high in the female placenta than the male placenta both in rodents and humans.^{23,24} In our EPS mouse model, placental *OGT* is further reduced by stress in both sexes, resulting in significant and lasting reductions throughout gestation in *OGT* levels in male placentas owing to the lower starting point of *OGT* in males.^{34,35}

Establishing causality with placental-specific gene targeting of *OGT*, we recapitulated the EPS phenotype, confirming the importance of placental *OGT* in neurodevelopmental programming and metabolic regulation.³⁴ To do this, we utilized a trophoblast-specific transgenic mouse to express Cre-recombinase (only in the placenta) and crossed this mouse with a conditional *OGT* mouse, in which *loxP* insertion around the *OGT* gene allows for excision of *OGT* in the presence of the Cre-recombinase, in order to examine the role of placental *OGT* in neurodevelopment, according to gene dosage.^{23,24} Hemizygous and homozygous deletion of *OGT* in trophoblast cells of the placenta completely recapitulated our EPS phenotype, producing offspring with significant hypothalamic programming changes, including dramatic and significant deficiency in mitochondrial function and transcriptional changes within the paraventricular nucleus of the hypothalamus (PVN), the energy and stress-regulating center of the brain.²³

To delve even deeper into the cellular mechanisms of the placenta that may relate to brain development and explain these sex differences, we need to understand the molecular role of *OGT* in this tissue. One major function of *OGT* is to broadly control transcriptional repression via its stabilization of a protein, enhancer of zeste homolog 2 (EZH2)—a histone H3K27 methyltransferase—thereby increasing levels of methylation of this global histone repressive mark, H3K27me3 (histone H3 trimethylated at lysine 27) (Figure 1).^{52,57,58} Therefore, from these data we can propose a hypothesis for a sex-specific mechanism that broadly controls pla-

cental transcription and function whereby the low levels of OGT in males result in low levels of the histone repressive mark, H3K27me3. This would position male chromatin to be in a much more “reactive” state, poised to produce robust transcriptional responses to maternal insults such as stress.^{21,23-25,27}

Within the placenta, other sex-specific cellular processes have been examined, with the most common being related to maternal stress and glucocorticoids and the glucocorticoid receptor and/or inactivating enzyme levels within the placenta.⁵⁹⁻⁶⁴ Within the placenta, maternal stress and glucocorticoids decrease placental glycogen levels in female, but not male, tissue suggesting important fetal sex differences in energy availability during maternal stress that may allow an adaptive response in females that is not found in males.^{20,65} Although we are beginning to identify placental cellular processes that may contribute to the sex bias in neurodevelopmental disorder risk, we still lack a clear understanding of what the important sex-specific signals are that ultimately reach and program the brain during gestation.

Transplacental signals

Most of what is reported for changes in placental signals related to neurodevelopmental disorders or sex differences is limited to associational studies, as it is extremely difficult to demonstrate direct placental transmission




Figure 1. Schematic illustrating the chromatin regulation and transcriptional control by O-linked N-acetylglucosamine transferase (OGT). For transcriptional silencing, OGT controls protein stability of EZH2, the H3K27me3 methyltransferase. Utx is a H3K27me3 demethylase. Increased OGT positively correlates with levels of H3K27me3 and transcriptional repression. EZH2, enhancer of zeste homolog 2; H3, histone H3; K27, lysine 27; Me, methyl; OGT, O-linked N-acetylglucosamine transferase; Utx, a histone H3-lysine 27 demethylase

in animal models or in humans due to the complexity in determining the origination source of any given molecule.^{20,24,32,55,66-74} However, one of the strongest stories we currently have centers on the production of, and changes in, fetal testosterone exposure and neurodevelopmental programming.^{13,20,23,68-75} The sexually dimorphic brain is organized during a critical window of development via hormone exposure, with males experiencing elevated testosterone levels during the process of normal testes development, largely in utero, although the exact timing of this is species dependent (as reviewed in reference 76). Aromatization of this testosterone to estradiol in the brain drives masculinization, an active process affecting cell differentiation and connectivity in the brain in most mammals. In humans, a strong correlation exists between fetal testosterone levels and neurodevelopmental disease risk, as well as with adult cognitive and behavioral stress reactivity, supporting a potential importance of these processes involved in establishing the sexually dimorphic brain, which occurs very early in brain development.^{68,69,73-75} Programming of important regulatory brain regions, including limbic regions such as the amygdala and hippocampus, and the neuroendocrine hypothalamus, via this developmental testosterone exposure and its effects on cell-migration patterns, probably also contributes to sex differences in how the individual responds to environmental challenges throughout life.⁷⁷

The current causal question then is how do sex differences within the placenta relate to disruptions in the sexually dimorphic brain? It is intriguing that whereas autism has been described as the “hypermale brain,” and many autism traits are directly correlated with high fetal testosterone levels,^{68,75} schizophrenia, especially in men, has been described as the “hypomale brain,” and studies in male patients have reported low levels of testosterone^{78,79} and reduced gestational programming by testosterone.^{80,81} Therefore, one mechanism that may place placental function at the root of sex biases in neurodevelopmental disease risk is that of variations in steroid hormone production during key periods of brain development.^{23,32,82} The placenta has robust and dynamic steroidogenic activity, and in many species, the male placenta produces high levels of testosterone during gestation.^{23,83-85}

As an example of placental steroid hormones influencing the brain, in our EPS model, we identified a significant reduction in OGT association with the

Brief report

17- β -hydroxysteroid dehydrogenase-3 (*Hsd17b3*) gene locus in male EPS placentas by ChIP-Seq analyses (chromatin immunoprecipitation combined with massively parallel DNA sequencing). As expected, this finding correlated with a significant reduction in Hsd17 β 3 expression in these placentas.²³ Hsd17 β 3 is a steroidogenic enzyme that converts androstenedione into testosterone. When we subsequently examined placental

hormones in EPS male placentas, we found a concordant significant reduction in testosterone and increase in androstenedione levels, supporting a functional outcome resulting from EPS exposure.²³ Importantly, adult EPS males also demonstrated a dysmasculinized phenotype, where their stress behaviors, cognitive function, and spatial strategies were much more similar to the control females than to the control males.^{39,56,86} We are certainly not the first to report such prenatal programming of dysmasculinized phenotypes due to maternal stress.⁸⁷⁻⁹²

Altogether, this Brief Report provides provocative discussion as to the potential critical involvement of the placenta in producing sex-specific responses to maternal insults in pregnancy that impact neurodevelopment and, ultimately, disease risk (Figure 2). Although steroid hormones may be involved as transplacental signals, these sex differences begin with genes from the sex chromosomes that drive sex-specific transcriptional regulation. A significant increase in the X-linked gene *OGT* and its control of the transcriptional repressive mark H3K27me3 in female (XX) compared with male (XY) trophoblast cells may position female placentas to be less responsive to a dynamic or changing gestational environment, providing a protective effect for the developing brain. Further studies are necessary to determine additional molecular mechanisms involved and their potential contribution to the known sex bias in neurodevelopmental disorders. \square




Figure 2. Summary illustration of the hypothesis regarding sex differences in chromatin regulation via O-linked N-acetylglucosamine transferase (OGT) and its regulation of the repressive histone mark H3K27me3 that may underlie the male bias for neurodevelopmental disorders. The increased expression of the X-linked gene *OGT*, in the female placenta provides them with a starting point from which a reduction due to a prenatal insult, such as maternal stress (orange lightning bolt), still affords significantly higher female *OGT* levels than found in males. Maternal insults can promote widespread transcriptional changes in male placentas (blue) but relatively little in female tissue due to differences in transcriptional repression (pink). The increased H3K27me3 levels in female placentas may then promote sex differences in transcriptional responses and, ultimately, placental function in response to early prenatal stress, placing males at an increased risk for neurodevelopmental disorders stemming from prenatal insults. EPS, early prenatal stress; H3K27me3, histone H3 trimethylated at lysine 27; Neurodev, neurodevelopmental disorder; OGT, O-linked N-acetylglucosamine transferase; XX, female; XY, male.

Conflict of Interest: The author declares no conflict of interest.

REFERENCES

1. Walker CK, Krakowiak P, Baker A, Hansen RL, Ozonoff S, Hertz-Picciotto I. Preeclampsia, placental insufficiency, and autism spectrum disorder or developmental delay. *JAMA Pediatr.* 2015;169(2):154-162.
2. Braithwaite EC, Kundakovic M, Ramchandani PG, Murphy SE, Champagne FA. Maternal prenatal depressive symptoms predict infant *NR3C1* 1F and *BDNF* IV DNA methylation. *Epigenetics.* 2015;10(5):408-417.
3. Togher KL, O'Keefe MM, Khashan AS, Gutierrez H, Kenny LC, O'Keefe GW. Epigenetic regulation of the placental HSD11B2 barrier and its role as a critical regulator of fetal development. *Epigenetics.* 2014;9(6):816-822.
4. Avishai-Eliner S, Brunson KL, Sandman CA, Baram TZ. Stressed-out, or in (utero)? *Trends Neurosci.* 2002;25(10):518-524.
5. Buss C, Entringer S, Davis EP, et al. Impaired executive function mediates the association between maternal pre-pregnancy body mass index and child ADHD symptoms. *PLoS One.* 2012;7(6):e37758.
6. Glynn LM, Wadhwa PD, Dunkel-Schetter C, Chicx-Demet A, Sandman CA. When stress happens matters: effects of earthquake timing on stress responsiveness in pregnancy. *Am J Obstet Gynecol.* 2001;184(4):637-642.
7. Sandman CA, Glynn LM, Davis EP. Is there a viability-vulnerability tradeoff? Sex differences in fetal programming. *J Psychosom Res.* 2013;75(4):327-335.

8. Wadhwa PD, Sandman CA, Garite TJ. The neurobiology of stress in human pregnancy: implications for prematurity and development of the fetal central nervous system. *Prog Brain Res.* 2001;133:131-142.
9. Wadhwa PD, Sandman CA, Porto M, Dunkel-Schetter C, Garite TJ. The association between prenatal stress and infant birth weight and gestational age at birth: a prospective investigation. *Am J Obstet Gynecol.* 1993;169(4):858-865.
10. Khashan AS, Abel KM, McNamee R, et al. Higher risk of offspring schizophrenia following antenatal maternal exposure to severe adverse life events. *Arch Gen Psychiatry.* 2008;65(2):146-152.
11. Sandman CA, Glynn LM, Davis EP. Is there a viability-vulnerability tradeoff? Sex differences in fetal programming. *J Psychosom Res.* 2013;75(4):327-335.
12. Buss C, Davis EP, Class QA, et al. Maturation of the human fetal startle response: evidence for sex-specific maturation of the human fetus. *Early Hum Dev.* 2009;85(10):633-638.
13. Salem JE, Kring AM. The role of gender differences in the reduction of etiologic heterogeneity in schizophrenia. *Clin Psychol Rev.* 1998;18(7):795-819.
14. Castle DJ, Wessely S, Murray RM. Sex and schizophrenia: effects of diagnostic stringency, and associations with and premorbid variables. *Br J Psychiatry.* 1993;162:658-664.
15. Castle DJ, Murray RM. The neurodevelopmental basis of sex differences in schizophrenia. *Psychol Med.* 1991;21(3):565-575.
16. Geschwind N, Galaburda AM. Cerebral lateralization. Biological mechanisms, associations, and pathology: III. A hypothesis and a program for research. *Arch Neurol.* 1985;42(7):634-654.
17. Geschwind N, Galaburda AM. Cerebral lateralization. Biological mechanisms, associations, and pathology: II. A hypothesis and a program for research. *Arch Neurol.* 1985;42(6):521-552.
18. Geschwind N, Galaburda AM. Cerebral lateralization. Biological mechanisms, associations, and pathology: I. A hypothesis and a program for research. *Arch Neurol.* 1985;42(5):428-459.
19. McGlone J. Sex differences in functional brain asymmetry. *Cortex.* 1978;14(1):122-128.
20. Davis EP, Pfaff D. Sexually dimorphic responses to early adversity: implications for affective problems and autism spectrum disorder. *Psychoneuroendocrinology.* 2014;49:11-25.
21. Nugent BM, Bale TL. The omniscient placenta: metabolic and epigenetic regulation of fetal programming. *Front Neuroendocrinol.* 2015;39:28-37.
22. Gabory A, Roseboom TJ, Moore T, Moore LG, Junien C. Placental contribution to the origins of sexual dimorphism in health and diseases: sex chromosomes and epigenetics. *Biol Sex Differ.* 2013;4(1):5.
23. Howerton CL, Bale TL. Targeted placental deletion of OGT recapitulates the prenatal stress phenotype including hypothalamic mitochondrial dysfunction. *Proc Natl Acad Sci U S A.* 2014;111(26):9639-9644.
24. Howerton CL, Morgan CP, Fischer DB, Bale TL. O-GlcNAc transferase (OGT) as a placental biomarker of maternal stress and reprogramming of CNS gene transcription in development. *Proc Natl Acad Sci U S A.* 2013;110(13):5169-5174.
25. Howerton CL, Bale TL. Prenatal programming: at the intersection of maternal stress and immune activation. *Horm Behav.* 2012;62(3):237-242.
26. Charil A, Laplante DP, Vaillancourt C, King S. Prenatal stress and brain development. *Brain Res Rev.* 2010;65(1):56-79.
27. Bronson SL, Bale TL. The placenta as a mediator of stress effects on neurodevelopmental reprogramming. *Neuropsychopharmacology.* 2016;41(1):207-218.
28. Bale TL. Sex differences in prenatal epigenetic programming of stress pathways. *Stress.* 2011;14(4):348-356.
29. Bronson SL, Bale TL. Prenatal stress-induced increases in placental inflammation and offspring hyperactivity are male-specific and ameliorated by maternal antiinflammatory treatment. *Endocrinology.* 2014;155(7):2635-2646.
30. Dunn GA, Morgan CP, Bale TL. Sex-specificity in transgenerational epigenetic programming. *Horm Behav.* 2011;59(3):290-295.
31. Gabory A, Roseboom TJ, Moore T, Moore LG, Junien C. Placental contribution to the origins of sexual dimorphism in health and diseases: sex chromosomes and epigenetics. *Biol Sex Differ.* 2013;4(1):5.
32. Goel N, Bale TL. Examining the intersection of sex and stress in modelling neuropsychiatric disorders. *J Neuroendocrinol.* 2009;21(4):415-420.
33. Graves JA. Review: Sex chromosome evolution and the expression of sex-specific genes in the placenta. *Placenta.* 2010;31(suppl):S27-S32.
34. Howerton CL, Bale TL. Targeted placental deletion of OGT recapitulates the prenatal stress phenotype including hypothalamic mitochondrial dysfunction. *Proc Natl Acad Sci U S A.* 2014;111(26):9639-9644.
35. Howerton CL, Morgan CP, Fischer DB, Bale TL. O-GlcNAc transferase (OGT) as a placental biomarker of maternal stress and reprogramming of CNS gene transcription in development. *Proc Natl Acad Sci U S A.* 2013;110(13):5169-5174.
36. Kim DR, Bale TL, Epperson CN. Prenatal programming of mental illness: current understanding of relationship and mechanisms. *Curr Psychiatry Rep.* 2015;17(2):5.
37. Mao J, Zhang X, Sieli PT, Falduto MT, Torres KE, Rosenfeld CS. Contrasting effects of different maternal diets on sexually dimorphic gene expression in the murine placenta. *Proc Natl Acad Sci U S A.* 2010;107(12):5557-5562.
38. McKendry AA, Palliser HK, Yates DM, Walker DW, Hirst JJ. The effect of betamethasone treatment on neuroactive steroid synthesis in a foetal Guinea pig model of growth restriction. *J Neuroendocrinol.* 2010;22(3):166-174.
39. Mueller BR, Bale TL. Sex-specific programming of offspring emotionality after stress early in pregnancy. *J Neurosci.* 2008;28(36):9055-9065.
40. Pankevich DE, Mueller BR, Brockel B, Bale TL. Prenatal stress programming of offspring feeding behavior and energy balance begins early in pregnancy. *Physiol Behav.* 2009;98(1-2):94-102.
41. Rich-Edwards J, Krieger N, Majzoub J, Zierler S, Lieberman E, Gillman M. Maternal experiences of racism and violence as predictors of preterm birth: rationale and study design. *Paediatr Perinat Epidemiol.* 2001;15(suppl 2):124-135.
42. Rivera RM, Stein P, Weaver JR, Mager J, Schultz RM, Bartolomei MS. Manipulations of mouse embryos prior to implantation result in aberrant expression of imprinted genes on day 9.5 of development. *Hum Mol Genet.* 2008;17(1):1-14.
43. Smith JT, Waddell BJ. Increased fetal glucocorticoid exposure delays puberty onset in postnatal life. *Endocrinology.* 2000;141(7):2422-2428.
44. Soares MJ, Chakraborty D, Karim Rumi MA, Konno T, Renaud SJ. Rat placentation: an experimental model for investigating the hemochorial maternal-fetal interface. *Placenta.* 2012;33(4):233-243.
45. Konno T, Rempel LA, Arroyo JA, Soares MJ. Pregnancy in the brown Norway rat: a model for investigating the genetics of placentation. *Biol Reprod.* 2007;76(4):709-718.
46. Rossant J, Cross JC. Placental development: lessons from mouse mutants. *Nat Rev Genet.* 2001;2(7):538-548.
47. Myatt L. Placental adaptive responses and fetal programming. *J Physiol.* 2006;572(pt 1):25-30.
48. Mao J, Zhang X, Sieli PT, Falduto MT, Torres KE, Rosenfeld CS. Contrasting effects of different maternal diets on sexually dimorphic gene expression in the murine placenta. *Proc Natl Acad Sci U S A.* 2010;107(12):5557-5562.
49. Keverne EB. Genomic imprinting and the evolution of sex differences in mammalian reproductive strategies. *Adv Genet.* 2007;59:217-243.
50. Keverne EB. Genomic imprinting, action, and interaction of maternal and fetal genomes. *Proc Natl Acad Sci U S A.* 2015;112(22):6834-6840.
51. Clifton VL. Review: Sex and the human placenta: mediating differential strategies of fetal growth and survival. *Placenta.* 2010;31(suppl):S33-S39.
52. Nugent BM, Bale TL. The omniscient placenta: metabolic and epigenetic regulation of fetal programming. *Front Neuroendocrinol.* 2015;39:28-37.
53. Bronson SL, Bale TL. The placenta as a mediator of stress effects on neurodevelopmental reprogramming. *Neuropsychopharmacology.* 2016;41(1):207-218.
54. Hsiao EY, Patterson PH. Activation of the maternal immune system induces endocrine changes in the placenta via IL-6. *Brain Behav Immun.* 2011;25(4):604-615.
55. Hsiao EY, Patterson PH. Placental regulation of maternal-fetal interactions and brain development. *Dev Neurobiol.* 2012;72(10):1317-1326.
56. Morgan CP, Bale TL. Early prenatal stress epigenetically programs dysmasculinization in second-generation offspring via the paternal lineage. *J Neurosci.* 2011;31(33):11748-11755.

Brief report

57. Chu CS, Lo PW, Yeh YH, et al. O-GlcNAcylation regulates EZH2 protein stability and function. *Proc Natl Acad Sci U S A*. 2014;111(4):1355-1360.
58. Ruan HB, Singh JP, Li MD, Wu J, Yang X. Cracking the O-GlcNAc code in metabolism. *Trends Endocrinol Metab*. 2013;24(6):301-309.
59. Cottrell EC, Seckl JR. Prenatal stress, glucocorticoids and the programming of adult disease. *Front Behav Neurosci*. 2009;3:19.
60. Drake AJ, Walker BR, Seckl JR. Intergenerational consequences of fetal programming by in utero exposure to glucocorticoids in rats. *Am J Physiol Regul Integr Comp Physiol*. 2005;288(1):R34-R38.
61. Holmes MC, Abrahamsen CT, French KL, Paterson JM, Mullins JJ, Seckl JR. The mother or the fetus? 11-hydroxysteroid dehydrogenase type 2 null mice provide evidence for direct fetal programming of behavior by endogenous glucocorticoids. *J Neurosci*. 2006;26(14):3840-3844.
62. Mairesse J, Lesage J, Breton C, et al. Maternal stress alters endocrine function of the feto-placental unit in rats. *Am J Physiol Endocrinol Metab*. 2007;292(6):E1526-E1533.
63. Nyirenda MJ, Seckl JR. Intrauterine events and the programming of adulthood disease: the role of fetal glucocorticoid exposure (Review). *Int J Mol Med*. 1998;2(5):607-614.
64. Seckl JR, Holmes MC. Mechanisms of disease: glucocorticoids, their placental metabolism and fetal 'programming' of adult pathophysiology. *Nat Clin Pract Endocrinol Metab*. 2007;3(6):479-488.
65. O'Connell BA, Moritz KM, Walker DW, Dickinson H. Treatment of pregnant spiny mice at mid gestation with a synthetic glucocorticoid has sex-dependent effects on placental glycogen stores. *Placenta*. 2013;34(10):932-940.
66. Gore AC, Martien KM, Gagnidze K, Pfaff D. Implications of prenatal steroid perturbations for neurodevelopment, behavior, and autism. *Endocr Rev*. 2014;35(6):961-991.
67. O'Donnell K, O'Connor TG, Glover V. Prenatal stress and neurodevelopment of the child: focus on the HPA axis and role of the placenta. *Dev Neurosci*. 2009;31(4):285-292.
68. Baron-Cohen S, Auyeung B, Norgaard-Pedersen B, et al. Elevated fetal steroidogenic activity in autism. *Mol Psychiatry*. 2015;20(3):369-376.
69. Chura LR, Lombardo MV, Ashwin E, et al. Organizational effects of fetal testosterone on human corpus callosum size and asymmetry. *Psychoneuroendocrinology*. 2010;35(1):122-132.
70. Knickmeyer R, Baron-Cohen S, Raggatt P, Taylor K. Foetal testosterone, social relationships, and restricted interests in children. *J Child Psychol Psychiatry*. 2005;46(2):198-210.
71. Knickmeyer R, Baron-Cohen S, Raggatt P, Taylor K, Hackett G. Fetal testosterone and empathy. *Horm Behav*. 2006;49(3):282-292.
72. Knickmeyer RC, Wheelwright S, Taylor K, Raggatt P, Hackett G, Baron-Cohen S. Gender-typed play and amniotic testosterone. *Dev Psychol*. 2005;41(3):517-528.
73. Lombardo MV, Ashwin E, Auyeung B, et al. Fetal programming effects of testosterone on the reward system and behavioral approach tendencies in humans. *Biol Psychiatry*. 2012;72(10):839-847.
74. Lombardo MV, Ashwin E, Auyeung B, et al. Fetal testosterone influences sexually dimorphic gray matter in the human brain. *J Neurosci*. 2012;32(2):674-680.
75. Baron-Cohen S, Lombardo MV, Auyeung B, Ashwin E, Chakrabarti B, Knickmeyer R. Why are autism spectrum conditions more prevalent in males? *PLoS Biol*. 2011;9(6):e1001081.
76. McCarthy MM. Multifaceted origins of sex differences in the brain. *Philos Trans R Soc Lond B Biol Sci*. 2016;371(1688):20150106.
77. Bale TL, Baram TZ, Brown AS, et al. Early life programming and neurodevelopmental disorders. *Biol Psychiatry*. 2010;68(4):314-319.
78. Goyal RO, Sagar R, Ammini AC, Khurana ML, Alias AG. Negative correlation between negative symptoms of schizophrenia and testosterone levels. *Ann N Y Acad Sci*. 2004;1032:291-294.
79. da Silva TL, Ravindran AV. Contribution of sex hormones to gender differences in schizophrenia: a review. *Asian J Psychiatr*. 2015;18:2-14.
80. Markham JA. Sex steroids and schizophrenia. *Rev Endocr Metab Disord*. 2012;13(3):187-207.
81. Collinson SL, Lim M, Chaw JH, et al. Increased ratio of 2nd to 4th digit (2D:4D) in schizophrenia. *Psychiatry Res*. 2010;176(1):8-12.
82. Sarkar P, Bergman K, O'Connor TG, Glover V. Maternal antenatal anxiety and amniotic fluid cortisol and testosterone: possible implications for foetal programming. *J Neuroendocrinol*. 2008;20(4):489-496.
83. Gopalakrishnan K, Mishra JS, Chinnathambi V, et al. Elevated testosterone reduces uterine blood flow, spiral artery elongation, and placental oxygenation in pregnant rats. *Hypertension*. 2016;67(3):630-639.
84. Sathishkumar K, Balakrishnan M, Chinnathambi V, Chauhan M, Hankins GD, Yallampalli C. Fetal sex-related dysregulation in testosterone production and their receptor expression in the human placenta with preeclampsia. *J Perinatol*. 2012;32(5):328-335.
85. Sathishkumar K, Elkins R, Chinnathambi V, Gao H, Hankins GD, Yallampalli C. Prenatal testosterone-induced fetal growth restriction is associated with down-regulation of rat placental amino acid transport. *Reprod Biol Endocrinol*. 2011;9:110.
86. Mueller BR, Bale TL. Early prenatal stress impact on coping strategies and learning performance is sex dependent. *Physiol Behav*. 2007;91(1):55-65.
87. Ward IL. Prenatal stress feminizes and demasculinizes the behavior of males. *Science*. 1972;175(17):82-84.
88. Ward IL, Stehm KE. Prenatal stress feminizes juvenile play patterns in male rats. *Physiol Behav*. 1991;50(3):601-605.
89. Kapoor A, Kostaki A, Janus C, Matthews SG. The effects of prenatal stress on learning in adult offspring is dependent on the timing of the stressor. *Behav Brain Res*. 2009;197(1):144-149.
90. Kapoor A, Leen J, Matthews SG. Molecular regulation of the hypothalamic-pituitary-adrenal axis in adult male guinea pigs after prenatal stress at different stages of gestation. *J Physiol*. 2008;586(pt 17):4317-4326.
91. Liu L, Li A, Matthews SG. Maternal glucocorticoid treatment programs HPA regulation in adult offspring: sex-specific effects. *Am J Physiol Endocrinol Metab*. 2001;280(5):E729-E739.
92. Owen D, Matthews SG. Glucocorticoids and sex-dependent development of brain glucocorticoid and mineralocorticoid receptors. *Endocrinology*. 2003;144(7):2775-2784.